

AGROCELHONE C INJECTABLE

COMPOSICIÓN Y PROPIEDADES

LIQUID INJECTABLE FUMIGANT AL FUNGICIDE-SOIL DISINFECTANT

CHLOROPICRIN 99.5% (99,5% P/P) P/V
SPECIFIC WEIGHT 1.65 KG/L



■ CROPS

VEGETABLES, FLOWERS AND FRUIT CROPS

DOSE

200-400 KG/Ha equiv. to 121.2 - 242.4 l/Ha



■ Safety period: **21 days.**

■ Method of use: Mechanised injection.

■ Soil preparation

- Pull up and remove previous crop.
- Cross-tilling depth: 40-50 cm for herbaceous crops, 60-80 cm (mole drains) for trees.
- Surface cross-tilling (rotovator).
- Soil humidity must be approximately 50 %.
- Treatment can be all-terrain, in alternating strips, or localised over the crop line.

■ In localised treatment on beds or crop lines the applied dose shall be in gr/m² of actual area.

■ Post-treatment tasks

- The exposure time of the films (time for the product to be active), is approximately 12-14 days
- In all-terrain treatments, once the films are removed, cross-till with shanks or cultivator to aerate the soil. In localised treatments, only the treated strip shall be tilled.

Recommendations

AGROCELHONE CE is recommended for controlling mainly soil fungi.