

AGROCELHONE F INJECTABLE

COMPOSITION AND PROPERTIES

LIQUID INJECTABLE FUMIGANT AL FUNGICIDE-NEMATICIDE-SOIL DISINFECTANT

DICHLOROPROPENE 39,5% (39.5% P/P)P/V

CHLOROPICRIN 59,5% (59.5% P/P) P/V

SPECIFIC WEIGHT 1.47 KG/L



■ CROPS

VEGETABLES, FLOWERS AND FRUIT CROPS

DOSE

350 -500 KG/Ha equiv. to 245-350 l/Ha



■ Safety period: **21 days.**

■ Method of use: Mechanised injection.

■ Soil preparation

- Pull up and remove previous crop.
- Cross-tilling depth: 40-50 cm for herbaceous crops, 60-80 cm (mole drains) for trees.
- Surface cross-tilling (rotovator).
- Soil humidity must be approximately 50 %.
- Treatment can be all-terrain, in alternating strips, or localised over the crop line.

■ In localised treatment on beds or crop lines the applied dose shall be in gr/m² of actual area.

■ Post-treatment tasks

- The exposure time of the films (time for the product to be active), is approximately 12-14 days
- In all-terrain treatments, once the films are removed, cross-till with shanks or cultivator to aerate the soil. In localised treatments, only the treated strip shall be tilled.

Recommendations

AGROCELHONE F is recommended when the main problem is soil fungi, even when other minor problems are present.